

1969. Generalized solitary waves in nonintegrable KdV equations

Zenonas Navickas¹, Minvydas Ragulskis², Romas Marcinkevicius³, Tadas Telksnys⁴

^{1, 2, 4}Department of Mathematical Modeling, Kaunas University of Technology, Kaunas, Lithuania

³Department of Software Engineering, Kaunas University of Technology, Kaunas, Lithuania

⁴Corresponding author

E-mail: ¹zenonas.navickas@ktu.lt, ²minvydas.ragulskis@ktu.lt, ³romas.marcinkevicius@ktu.lt,

⁴tadas.telksnys@ktu.lt

(Received 29 December 2015; received in revised form 17 February 2016; accepted 2 March 2016)

Abstract. The generalization of the classical Korteweg-de-Vries (KdV) solitary wave solution is presented in this paper. The amplitude and the propagation speed of generalized KdV solitary waves vary in time. Generating partial differential equations and conditions of existence of the generalized KdV solitary waves are derived using the inverse balancing method. Computational experiments illustrate the variety of new solitary solutions and their generating equations.

Keywords: Korteweg-de-Vries equation, solitary solution, existence condition.

1. Introduction

Solitary solutions of nonlinear partial differential equations play a major part in mathematical physics [1]. While most commonly solitary waves maintain constant amplitude and propagation speed (because they are obtained via the traveling wave substitution $\xi = \lambda x + \mu t$) recent developments include solitary solutions which have either a variable amplitude or propagation speed, or both. It is demonstrated in [2] that soliton's amplitude and width can alter during the propagation in the model of (2+1)-dimensional coupled nonlinear Schrodinger equations in a graded-index waveguide. It is shown that the maximum amplitude of the solitary solution to the (1+1)D CQNLS equation in the self-focusing quintic case does depend on the propagation distance [3].

A number of oscillating solitons such as the dromion and solitoff have been shown to satisfy the nonlinear (2+1)-dimensional Broer-Kaup (BK) equations in [4]. The deformed soliton, breather and rogue wave solutions of the inhomogeneous nonlinear Hirota equation with linear inhomogeneous coefficients and higher-order dispersion are discussed in [5]. The variational approach is employed in [6] to construct three classes approximate azimuthally modulated vortex solitons for the (2+1)-dimensional nonlinear Schrodinger equation with transverse nonperiodic modulation. Higher-order Dirac solitons, which have amplitude profiles that repeat periodically over large propagation distances are investigated in [7].

Solitary solutions are often considered in the context of birefringent optical fibers. In [8], an analytic model is constructed to demonstrate polarized soliton oscillations in a lossy elliptically low birefringent optical fiber. A study on the propagation of optical solitons through birefringent fibers in the presence of spatio-temporal dispersion is presented in [9]. The Riccati equation and Jacobian elliptic equation methods are used to study the propagation of optical solitons in birefringent fibers with parabolic law nonlinearity [10]. Bright and dark 1-soliton solutions of systems modeling birefringent fibers are considered using several perturbation terms in [11]. The dynamics of soliton propagation through optical metamaterials can be found in [12].

There have also been reports of experimental evidence of solitary solutions with non-constant speed and amplitude. A discussion on the way spiralling elliptic solitons can be observed experimentally can be found in [13]. A study on soliton frequency shifts in few-cycle ultrashort laser pulses propagating through resonant media has recently been published in [14]; self-decelerating Airy-Bessel light bullets are discussed in [15].

Recent exact solitary solution derivation techniques have demonstrated that the field is no longer limited to the classical soliton. The curved soliton solution $w = \operatorname{sech}^2(x + \exp(-t))$ to

the damped externally excited KdV equation has been obtained in [16]; the embedded and annula soliton solutions to the (2+1)-dimensional Burgers equation have been discussed in [17].

The main objective of this article is to provide an analytical framework for the determination of the generating equations which possess generalized solitary solutions with non-constant amplitude and propagation speed. The inverse balancing method is used to identify classes of nonlinear partial differential equations that admit such solitary solutions. Conditions of existence for generalized solitary solutions are derived; computational experiments illustrate the variety of new solitary solutions and their generating equations.

2. Korteweg-de-Vries solitary waves

It is well-known that the Korteweg-de-Vries (KdV) equation:

$$w_t - 6ww_x + w_{xxx}''' = 0, \quad (1)$$

which models one-dimensional shallow water waves with small amplitudes does admit the solitary-wave solution:

$$w = w(t, x) = \gamma + \frac{\alpha \exp(x - st)}{(1 + \exp(x - st))^2}, \quad (2)$$

where $\gamma, \alpha, s \in \mathbb{R}$ are parameters that can be computed from Eq. (1) [1]. Note that s corresponds to the constant propagation speed of the wave; α corresponds to the amplitude and γ is the level of undisturbed water. The plot of Eq. (2) is given in Fig. 1.

In this paper, the following generalization of Eq. (2) is considered:

$$w = w(t, x) = \gamma + \frac{u(t) \exp(v(t, x))}{(1 + \exp(v(t, x)))^2}, \quad (3)$$

where $u(t)$ corresponds to the variable amplitude of the solitary wave and $v(t, x)$ to the propagation speed (which need not be constant). The functions u, v are continuous, differentiable and do not have essential singularities. This generalization allows the consideration of waves with soliton-like characteristics – but with variable speed and amplitude.




Fig. 1. Classical KdV solitary wave Eq. (2) with $\alpha = s = 1$

3. Generating equations of generalized KdV solitary waves

Partial differential equations that admit the solution Eq. (3) are derived in this section. Conditions of the existence of solutions Eq. (3) in a class of differential equations are considered.

3.1. Derivation of generating PDEs

Eq. (3) can be rewritten as:

$$w(t, x) = \tilde{w}(u, v) = \gamma + \frac{u \exp(v)}{(1 + \exp(v))^2}. \quad (4)$$

Differentiating Eq. (4) with respect to u, v yields:

$$\tilde{w}'_u = \frac{\exp(v)}{(1 + \exp(v))^2} = \frac{\tilde{w} - \gamma}{u}, \quad \tilde{w}'_v = \frac{u \exp(v) (\exp(v) - 1)}{(1 + \exp(v))^3} = \frac{\exp(v) - 1}{\exp(v) + 1} \tilde{w}. \quad (5)$$

The inverse balancing can now be used: considering Eqs. (4), (5) algebraic equations, the variables u and $\exp(v)$ can be expressed as:

$$u = \frac{\tilde{w} - \gamma}{\tilde{w}'_u}, \quad \exp(v) = \frac{\tilde{w} + \tilde{w}'_v}{\tilde{w} - \tilde{w}'_v}. \quad (6)$$

Inserting Eq. (6) into Eq. (4) yields the partial differential equation:

$$(\tilde{w}'_v)^2 = (1 - 4\tilde{w}'_u)(\tilde{w} - \gamma)^2. \quad (7)$$

Returning to the original variables yields:

$$\tilde{w}'_v = \frac{w'_x}{v'_x}, \quad \tilde{w}'_u = \frac{v'_x w'_t - v'_t w'_x}{u'_t v'_x}. \quad (8)$$

Thus Eq. (7) can be rewritten as:

$$(w'_x)^2 - \frac{4v'_x v'_t}{u'_t} w'_x (w - \gamma)^2 + \frac{4(v'_x)^2}{u'_t} w'_t (w - \gamma)^2 = (v'_x)^2 (w - \gamma)^2. \quad (9)$$

Note that the generating equations of the classical KdV solitary solution Eq. (2) are special cases of Eqs. (7) and (9) when $u'_t \rightarrow 0, v'_x \rightarrow 1, v'_t \rightarrow -s$:

$$(w'_x)^2 = (4w - \gamma - 1)(w - \gamma)^2, \quad (w'_t)^2 = s(4w - \gamma - 1)(w - \gamma)^2. \quad (10)$$

Both equations defined in Eq. (10) have solution Eq. (2). Subsequent differentiation and use of Eq. (10) yields the KdV Eq. (1).

The equations (9) are not integrable; however, it has been shown that solitary solutions exist in many nonintegrable systems [18]. Furthermore, their properties can be different than those that are encountered in integrable systems, including non-constant amplitude and propagation speed [18].

3.2. Derivation of generalized solitary solutions and their conditions

Suppose the following partial differential equation is given:

$$(w'_x)^2 + A_2(t, x)(w - \gamma)^2 w'_x + A_1(t, x)(w - \gamma)^2 w'_t = A_0(t, x)(w - \gamma)^2, \quad (11)$$

where A_2, A_1, A_0 are known functions of t, x . Comparing Eqs. (9) and (11) yields that Eq. (11) admits the generalized KdV solitary solution Eq. (3) if the following equations hold:

$$A_2(t, x) = -\frac{4v'_t v'_x}{u'_t}, \quad (12)$$

$$A_1(t, x) = \frac{4(v'_x)^2}{u'_t}, \quad (13)$$

$$A_0(t, x) = (v'_x)^2. \quad (14)$$

Eqs. (13), (14) yield:

$$u'_t = \frac{4A_0(t, x)}{A_1(t, x)}. \quad (15)$$

Let us introduce the function B :

$$B(t) = \frac{4A_0(t, x)}{A_1(t, x)}. \quad (16)$$

Note that B depends only on t , because $u = u(t)$.

The consistency condition $(v'_t)'_x = (v'_x)'_t$ and Eqs. (12)-(14) results in the constraint:

$$\begin{aligned} -A_1^2(t, x)(A_0(t, x))'_t &= 2A_0(t, x) \left(A_1(t, x)(A_2(t, x))'_x - A_2(t, x)(A_1(t, x))'_x \right) \\ &+ A_2(t, x)A_1(t, x)(A_0(t, x))'_x. \end{aligned} \quad (17)$$

Furthermore, if only real-valued solutions are considered, the condition:

$$A_0(t, x) \geq 0, \quad t, x \in \mathbb{R}, \quad (18)$$

must be satisfied. Then Eqs. (13), (14) have the solution:

$$v'_t = \pm \frac{A_2(t, x)\sqrt{A_0(t, x)}}{A_1(t, x)}, \quad v'_x = \mp \sqrt{A_0(t, x)}. \quad (19)$$

From Eqs. (15) and (19) it follows that the parameters of the generalized KdV solitary wave read:

$$u = \int_{t_0}^t B(\tau) d\tau + u_0, \quad (20)$$

$$v = \pm \left(\int_{x_0}^x \sqrt{A_0(t, \xi)} d\xi - \int_{t_0}^t \frac{A_2(\tau, x_0)\sqrt{A_0(\tau, x_0)}}{A_1(\tau, x_0)} d\tau \right). \quad (21)$$

Note that:

$$w = \gamma + \frac{u(t) \exp(v(t, x))}{(1 + \exp(v(t, x)))^2} = + \frac{u(t) \exp(-v(t, x))}{(1 + \exp(-v(t, x)))^2}, \quad (22)$$

for all $v(t, x)$, thus only the positive-sign branch of Eq. (21) is considered further in this paper.

3.3. Cauchy initial value problem on Eq. (11)

The Cauchy initial value problem on the partial differential Eq. (11) can be posed in the

following way: given the Eq. (11), find a generalized KdV solitary solution Eq. (3) subject to initial conditions:

$$u(t_0) = u_0, \quad v(t_0, x_0) = v_0, \quad u_0, v_0 \in \mathbb{R}. \quad (23)$$

The derivations of Subsection 3.2 indicate the necessary and sufficient conditions for the solution to the discussed Cauchy problem. The partial differential Eq. (11) admits the generalized KdV solitary solution Eq. (3) if and only if the coefficients of the equation $A_2(t, x)$, $A_1(t, x)$, $A_0(t, x)$ satisfy the conditions Eqs. (16), (17) and (18).




Fig. 2. Decaying amplitude generalized KdV solitary wave Eq. (30) with $t_0 = 0$, $u_0 = 1$, $x_0 = v_0 = 0$

4. Several examples of generalized solitary solutions

4.1. Decaying amplitude solitary wave

Suppose the following partial differential equation is given:

$$(w'_x)^2 - \frac{2 \exp(t^2)}{t} w^2 w'_x - \frac{2 \exp(t^2)}{t} w^2 w'_t = w^2. \quad (24)$$

The coefficients of the equation read:

$$A_2(t, x) = A_1(t, x) = -\frac{2 \exp(t^2)}{t}, \quad A_0(t, x) = 1, \quad B(t) = -2t \exp(t^2). \quad (25)$$

Eq. (24) has a generalized KdV solitary solution because conditions Eqs. (16), (17) and (18) hold true. Eqs. (15) and (19) read:

$$u'_t = 2t \exp(t^2), \quad (26)$$

$$v'_t = 1, \quad v'_x = 1. \quad (27)$$

The solution of Eqs. (26) and (27), according to Eqs. (20) and (21) reads:

$$u = \exp(-t^2) - \exp(-t_0^2) + u_0, \quad u_0 \in \mathbb{R}, \quad (28)$$

$$v = x - t + x_0 - t_0 + v_0, \quad v_0 \in \mathbb{R}. \quad (29)$$

Thus, the equation has the generalized solitary solution:

$$w = \frac{(\exp(-t^2) - \exp(-t_0^2) + u_0) \exp(x - t + x_0 - t_0 + v_0)}{(1 + \exp(x - t + x_0 - t_0 + v_0))^2}. \quad (30)$$

Note that the constants t_0, x_0 only determine the location of the peak amplitude of the solitary wave. The plot of Eq. (30) is given in Fig. 2. The propagation speed of the solitary wave remains constant throughout all values of $t \in \mathbb{R}$ – but while the amplitude of the solitary solution decays in time.

4.2. Decaying amplitude solitary wave with variable speed

Suppose the following partial differential equation is given:

$$(w'_x)^2 + 2 \exp(-t) \frac{(t^2 + 1)^2}{t} w^2 w'_x - 2 \frac{(t^2 + 1)^2}{t} w^2 w'_t = w^2. \quad (31)$$

Eq. (31) together with Eqs. (15) and (19) yields:

$$u'_t = -\frac{2t}{(t^2 + 1)^2}, \quad (32)$$

$$v'_t = \exp(-t), \quad v'_x = 1. \quad (33)$$




Fig. 3. Generalized KdV solitary wave with decaying amplitude and variable speed Eq. (36) with $t_0 = 0, u_0 = 1/10, x_0 = 0, v_0 = 1$

According to Eqs. (20) and (21) integration results in the functions:

$$u = \frac{t_0^2 - t^2}{(t_0^2 + 1)(t^2 + 1)} + u_0, \quad (34)$$

$$v = x - \exp(-t) + x_0 - \exp(-t_0) + v_0, \quad (35)$$

that yield the generalized solitary solution:

$$w = \frac{\left(\frac{t_0^2 - t^2}{(t_0^2 + 1)(t^2 + 1)} + u_0 \right) \exp(x - \exp(-t) + x_0 - \exp(-t_0) + v_0)}{(1 + \exp(x - \exp(-t) + x_0 - \exp(-t_0) + v_0))^2}. \quad (36)$$

The plot of Eq. (36) is given in Fig. 3. Note that this solitary solution has decaying amplitude and a variable propagation speed.

4.3. Solitary wave with bounded growing amplitude

Let the following partial differential equation be given:

$$(w'_x)^2 - \frac{6(t^4 + 1)}{t} w^2 w'_x + \frac{18(t^4 + 1)}{t} w^2 w'_t = 9w^2. \quad (37)$$

Following the same steps as in the previous examples, it is obtained that:

$$u'_t = \frac{2t}{t^4 + 1}, \quad (38)$$

$$v'_t = -1, \quad v'_x = 3. \quad (39)$$

Integrating yields the functions:

$$u = \arctan(t^2) - \arctan(t_0^2) + u_0, \quad (40)$$

$$v = 3x + t - 3x_0 - t_0 + v_0. \quad (41)$$

The solitary wave reads:

$$w = \frac{(\arctan(t^2) - \arctan(t_0^2) + u_0) \exp(3x + t - 3x_0 - t_0 + v_0)}{(1 + \exp(3x + t - 3x_0 - t_0 + v_0))^2}. \quad (42)$$

This type of generalized solitary solution has a growing amplitude as t increases (or decreases) from t_0 (Fig. 4). However, the amplitude of the solitary wave does not increase unboundedly.




Fig. 4. Generalized KdV solitary wave with bounded and growing amplitude Eq. (42) with $t_0 = x_0 = u_0 = v_0 = 0$

4.4. Breathers

4.4.1. Breather with constant amplitude

Suppose the following partial differential equation is given:

$$(w'_x)^2 - 8 \cot(t) w^2 w'_x + 16 \csc t w^2 w'_t = 4w^2. \quad (43)$$

In this case, Eqs. (15) and (19) read:

$$u'_t = -\sin t, \quad (44)$$

$$v'_t = \cos t, \quad v'_x = 2. \quad (45)$$

Integrating Eqs. (44) and (45) yields the solitary solution:

$$w = \frac{(\cos t - \cos t_0 + u_0) \exp(2x - \sin t + 2x_0 - \sin t_0 + v_0)}{(1 + \exp(2x - \sin t + 2x_0 - \sin t_0 + v_0))^2}. \quad (46)$$

Nonlinear waves such as Eq. (46), in which energy concentrates in an oscillatory fashion are called breathers [1]. The solution Eq. (46) is an example of a breather with constant amplitude: the amplitude of the wave varies for different t , but it does not decay (Fig. 5).




Fig. 5. Standing breather Eq. (46) with $t_0 = 0$, $x_0 = u_0 = 1$, $v_0 = 2$

4.4.2. Breather with decaying amplitude

Let the following partial differential equation be given:

$$(w'_x)^2 + \frac{4t^2 \sin t}{\sin t - \cos t} w^2 w'_x - \frac{4t^2}{\sin t - \cos t} w^2 w'_t = w^2. \quad (47)$$

Eqs. (15) and (19) read:

$$u'_t = \frac{t \cos t - \sin t}{t^2}, \quad (48)$$

$$v'_t = \sin t, \quad v'_x = 1. \quad (49)$$

Integration of Eqs. (48) and (49) results in the solitary solution:

$$w = \frac{\left(\frac{\sin t}{t} - \frac{\sin t_0}{t_0} + u_0\right) \exp(x - \cos t - x_0 + \cos t_0 + v_0)}{(1 + \exp(x - \cos t - x_0 + \cos t_0 + v_0))^2}. \quad (50)$$

The obtained solution Eq. (50) is a type of breather which has decaying amplitude (Fig. 6).

5. Conclusions

It is shown that the classical Korteweg-de-Vries solitary wave can be generalized to represent solitary wave structures with variable amplitude and propagation speed. The generating equations that have generalized KdV solitary solutions are derived using the inverse balancing method. The solution of this inverse problem helps to identify the form of partial differential equations that admit generalized KdV solitary solutions. Conditions of existence of such solutions in the obtained class of partial differential equations are derived.




Fig. 6. Standing breather with decaying amplitude Eq. (50) with $t_0 = 0$, $u_0 = x_0 = 1$, $v_0 = 0$

This technique of classical solitary wave generalization demonstrates that recent observations of solitary solutions with time-variable amplitude and propagation speed can be analysed by means of a lower order generating equation. This provides a solid foundation for the analysis of such solitary waves and provides the link between the parameters of the solution and the differential equation.

The application of the methods described in this paper to other forms of generalized solitary solutions remains a definite object of future research.

Acknowledgements

This research was funded by a Grant No. MIP078/15 from the Research Council of Lithuania.

References

- [1] Scott A. Encyclopedia of Nonlinear Science. Routledge, New York, 2004.
- [2] Xie X.-Y., Tian B., Sun W.-R., Sun Y. Bright solitons for the (2+1)-dimensional coupled nonlinear Schrodinger equations in a graded-index waveguide. Communications in Nonlinear Science and Numerical Simulation, Vol. 29, 2015, p. 300-306.
- [3] Goksel I., Antar N., Bakirtas I. Solitons of (1+1)D cubic-quintic nonlinear Schrodinger equation with pt-symmetric potentials. Optics Communications, Vol. 354, 2015, p. 277-285.
- [4] Pang Q. Study on the behavior of oscillating solitons using the (2+1)-dimensional nonlinear system. Applied Mathematics and Computation, Vol. 217, 2013, p. 2010-2015.
- [5] Liu X., Yong X., Huang Y., Yu R., Gaom J. Deformed soliton, breather and rogue wave solutions of an inhomogeneous nonlinear Hirota equation. Communications in Nonlinear Science and Numerical Simulation, Vol. 29, 2015, p. 257-266.
- [6] Lai X.-J., Dai C.-Q., Cai X.-O., Zhang J.-F. Azimuthally modulated vortex solitons in bulk dielectric media with a Gaussian barrier. Optics Communications, Vol. 353, 2015, p. 101-108.
- [7] Tran T. X., Duong D. C. Higher-order Dirac solitons in binary waveguide arrays. Annals of Physics, Vol. 361, 2015, p. 501-508.
- [8] Mahmood M. F. Analytic method for solving coupled nonlinear Schroedinger equations with oscillating terms describing polarized soliton oscillations. Mathematical and Computer Modeling, Vol. 36, 2002, p. 1259-1263.
- [9] Bhrawy A. H., Alshaery A. A., Hilal E. M., Savescu M., Milovic D., Khan K. R., Mahmood M. F., Jovanoski Z., Biswas A. Optical solitons in birefringent fibers with spatio-temporal dispersion. Optik – International Journal for Light and Electron Optics, Vol. 125, 2014, p. 4395-4944.
- [10] Zhou Q., Zhu Q., Biswas A. Optical solitons in birefringent fibers with parabolic law nonlinearity. Optica Applicata, Vol. 41, 2014, p. 399-409.
- [11] Biswas A., Khan K., Rahman A., Yildirim A., Hayat T., Aldossary O. M. Bright and dark optical solitons in birefringent fibers with Hamiltonian perturbations and Kerr law nonlinearity. Journal of Optoelectronics and Advanced Materials, Vol. 14, 2012, p. 571-576.

- [12] Xu Y., Savescu M., Khan K. R., Mahmood M. F., Biswas A., Belic M. Soliton propagation through nanoscale wave-guides in optical metamaterials. *Optics Laser Technology*, Vol. 77, 2016, p. 177-186.
- [13] Liang G., Li H. Polarized vector spiraling elliptic solitons in nonlocal nonlinear media. *Optics Communications*, Vol. 352, 2015, p. 39-44.
- [14] Song X., Yan M., Wu M., Sheng Z., Hao Z., Huang C., Yang W. Soliton frequency shifts in subwavelength structures. *Journal of Optics*, Vol. 17, 2015, p. 055503.
- [15] Zhong W.-P., Belic M., Zhang Y. Self-decelerating Airy-Bessel light bullets. *Journal of Physics B*, Vol. 48, Issue 17, 2015, p. 175401.
- [16] Gandarias M. L., Khalique C. M. Symmetries, solutions and conservation laws of a class of nonlinear dispersive wave equations. *Communications in Nonlinear Science and Numerical Simulation*, Vol. 32, 2016, p. 114-121.
- [17] Yang L., Du X., Yang Q. New variable separation solutions to the (2+1)-dimensional Burgers equation. *Applied Mathematics and Computation*, Vol. 273, 2015, p. 1271-1275.
- [18] Yang J. *Nonlinear waves in integrable and nonintegrable systems*. SIAM, Philadelphia, 2010.



Zenonas Navickas is a Professor of Mathematics at the Department of Mathematical Modeling, Kaunas University of Technology. He received his doctoral degree in 1975 from Vilnius University, Lithuania. His research interests include algebraic theory of operator analysis and its applications in science, technology and biomedicine.



Minvydas Ragulskis is a Professor of Mathematics at the Department of Mathematical Modeling, Kaunas University of Technology and is also the head of the Research Group for Mathematical and Numerical Analysis of Dynamical Systems. He received his Ph.D. from Kaunas University of Technology in 1992. His research areas are nonlinear dynamical systems and numerical analysis.



Romas Marcinkevicius is an Associate Professor at the Department of Software Engineering, Kaunas University of Technology. He received his Ph.D. from Kaunas University of Technology in 1983. His research areas are parallel computing and symbolic computation.



Tadas Telksnys received his M.Sc. in Mathematics from Kaunas University of Technology in 2015. He is currently a Ph.D. student at the Department of Mathematical Modeling, Kaunas University of Technology. His research interests include analytical solutions of nonlinear ordinary differential equations and linear recurring sequences.